

Traits	Ma (<i>n</i> = 9)	Ma × Du (<i>n</i> = 9)	LW (<i>n</i> = 10)	ZS (<i>n</i> = 10)	Pw (<i>n</i> = 10)	SEM	<i>p</i> value
Slaughter weight (kg)	100.7 ^b	108.4 ^a	100.9 ^b	103.3 ^{a,b}	102.0 ^b	0.780	0.008
Carcass length (cm)	71.6 ^d	80.9 ^c	97.7 ^a	86.0 ^b	83.1 ^{b,c}	1.309	< 0.001
Hot carcass weight (kg)	81.1	84.1	80.8	82.7	81.9	0.595	0.443
Slaughter yield (%)	80.5 ^a	77.5 ^b	80.1 ^a	80.0 ^a	80.3 ^a	0.230	< 0.001
Cold carcass weight (kg)	79.9	82.1	79.1	81.4	80.3	0.564	0.407
Chilling loss (%)	1.1	1.2	1	0.8	0.9	0.056	0.120
Loin eye area (mm ²)	26.1 ^c	40.0 ^b	48.0 ^a	37.8 ^b	37.9 ^b	1.217	< 0.001
Back-fat thickness (cm)	5.7 ^a	4.1 ^b	1.7 ^d	2.9 ^c	3.1 ^c	0.211	< 0.001

Ma: Mangalitsa, Ma × Du: crossbred Mangalitsa × Duroc, LW: Large White, ZS: Zlotnicka spotted, Pw: Pulawska; ^{a-d} different letters in the same row indicate significant statistical differences (Tukey's test *P* < 0.05), SEM: standard deviation of the mean.